



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 214 886 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
19.06.2002 Bulletin 2002/25

(51) Int Cl.7: **A23D 7/00, A61K 7/48,
A23L 1/24, A61K 47/14**

(21) Application number: **01129299.2**

(22) Date of filing: **13.12.2001**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

- **Asou, Yoshihide**
Sumida-ku, Tokyo 131-8501 (JP)
- **Kawai, Shigeru**
Sumida-ku, Tokyo 131-8501 (JP)
- **Nakajima, Yoshinobu**
Sumida-ku, Tokyo 131-8501 (JP)

(30) Priority: **15.12.2000 JP 2000381596**

(71) Applicant: **Kao Corporation**
Tokyo 103-8210 (JP)

(74) Representative: **HOFFMANN - EITLE**
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

(72) Inventors:
• **Shiiba, Daisuke**
Sumida-ku, Tokyo 131-8501 (JP)

(54) **Acidic oil-in-water type emulsion composition**

(57) The invention provides an acidic oil-in-water type emulsion composition which has an oil phase containing at least 20 % by weight of diacylglycerol and 0.5 to 5.0 % by weight of a crystallization inhibitor, and has excellent shelf stability at low temperatures though it

contains diacylglycerol at a high concentration, also good in appearance and flavor and useful as a diet or food for improving lipid metabolism.

EP 1 214 886 A1

Description

BACKGROUND OF THE INVENTION

Field of the Invention:

[0001] The present invention relates to an acidic oil-in-water type emulsion composition such as mayonnaise or dressing, which is excellent in emulsion stability even under low-temperature conditions such as in side refrigerators, good in appearance and flavor and useful as diet or food for improving lipid metabolism.

Description of the Background Art:

[0002] In recent years, it has been clarified that diacylglycerol (hereafter referred to as "DAG") has an obesity-preventing effect, an effect to prevent increase in weight, etc. (Japanese Patent Application Laid-Open No. 300828/1992, etc.), and it is attempted to incorporate this into various kinds of foods (Japanese Patent Application Laid-Open Nos. 8431/1991, 79858/1992 and 176181/1998, etc.). Furthermore, oil-in water type emulsions containing DAG and lecithin in an oil phase and protein in a water phase have been proposed (Japanese Patent Application Laid-Open No. 91451/1991)

[0003] In the case of, particularly, an acidic oil-in-water type emulsion composition such as mayonnaise or dressing, however, the low-temperature resistance has been insufficient in that a part of DAG in raw oil crystallizes under in-refrigerator conditions (-5 to 5°C) that cause demulsification (oil-off), and for mayonnaise, cracks were observed.

SUMMARY OF THE INVENTION

[0004] It is therefore an object of the present invention to provide an acidic oil-in-water type emulsion composition which has excellent shelf stability at low temperatures though containing DAG at a high concentration, also good in appearance and flavor and useful as a diet or food for improving lipid metabolism.

[0005] The present inventors have found that an acidic oil-in-water type emulsion composition satisfying the above requirements can be provided by containing a specified amount of crystallization inhibitors in an oil phase containing DAG.

[0006] Thus, according to the present invention, we inventors have provided an acidic oil-in-water type emulsion composition comprising an oil phase containing at least 20 % by weight of diacylglycerol and 0.5 to 5.0 % by weight of a crystallization inhibitor.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0007] DAG used in the present invention is obtained by esterifying hydroxyl groups at 1- and 2-positions or 1-and 3-positions of glycerol with fatty acids. The number of carbon atoms in fatty acid residues are preferably (but not limited to) 8 to 24, particularly 16 to 22. The amount of unsaturated fatty acid residues is preferably at least 55%, more preferably at least 70%, most preferably at least 90% based on all the fatty acid residue. DAG is obtained by any known process such as ester exchange reaction of vegetable oil and/or animal oil with glycerol or esterification reaction of fatty acids derived from the above oil with glycerol. The reaction method thereof may be either a chemical reaction by the aid of alkali catalysts, for example, or a biochemical reaction method using oil and/or fat hydrolase such as lipase. One or blends of such DAGs may be used. The content of DAG in the oil phase must be at least 20 % by weight, preferably at least 35 % by weight from the viewpoints of effectiveness as a diet and profitability. The oil phase may contain triacylglycerol, monoacylglycerol, fatty acids and the like in addition to DAG. These may be derived from any of vegetable oils and/or animal oils.

[0008] Examples of raw oil include vegetable oils such as soybean oil, rapeseed oil, sunflower oil, cotton-seed oil, corn oil, safflower oil, linseed oil, olive oil, rice oil and palm oil; animal oils such as beef tallow, lard and fish oil; which may be further fractionated or transesterified.

[0009] The oil phase content in the acidic oil-in-water type emulsion composition is preferably 5 to 85 % by weight, more preferably 10 to 80 % by weight, particularly preferably 20 to 80 % by weight. In particular, the oil phase in the mayonnaise-type composition is preferably 20 to 85 % by weight, more preferably 25 to 80 % by weight, particularly preferably 30 to 80 % by weight. The oil phase content in the dressing-type composition is preferably 5 to 60 % by weight, more preferably 10 to 55 % by weight, most preferably 20 to 50 % by weight.

[0010] The crystallization-inhibitor used in the present invention is preferably selected from polyglycerol fatty acid esters, sucrose fatty acid esters and sorbitan fatty acid esters. The polyglycerol fatty acid esters are preferably those in which the average polymerization degree of glycerol is 2 to 12, the number of carbon atoms in the fatty acid moiety

is 12 to 22, and the degree of esterification is at least 70 %. Furthermore, the HLB of polyglycerol fatty acid esters is preferably lower than 4.5, most preferably lower than 3.5. In the present invention, the average polymerization degree (Mw) is determined by the following equation from the relationship between the polymerization degree (n) and hydroxyl value (OHV) of polyglycerol and the theoretical value.

$$\text{OHV} = 56110(n + 2)/\text{Mw}$$

$$\text{Mw} = 74n + 18$$

[0011] The sucrose fatty acid esters are preferably those in which the degree of esterification with fatty acids having 12 to 22 carbon atoms is at least 50 %, and the remaining hydroxyl groups (not esterified with above-mentioned fatty acids) are acetylated. Furthermore, the sucrose fatty acid esters preferably have an HLB lower than 3, more preferably lower than 2. The sorbitan fatty acid esters are preferably those in which the number of carbon atoms in the fatty acid moiety is 12 to 22, and the HLB is lower than 3, more preferably lower than 2.5. One or more of these crystallization inhibitors may be used, and the content thereof must be 0.5 to 5.0 % by weight, preferably 0.6 to 3.0 % by weight from the viewpoints of sufficient crystallization-inhibiting effect at low temperatures and flavor as food. The polyglycerol fatty acid esters are particularly preferred from the viewpoint of crystallization-inhibiting effect at varied temperatures.

[0012] The water phase in the acidic oil-in-water type emulsion composition according to the present invention may contain water; vinegar; common salt; condiments such as sodium glutamate; saccharides such as sugar and starch syrup; seasonings such as *sake* and sweet *sake*; various kinds of vitamins; organic acids; spices; various kinds of vegetables or fruits; thickeners such as xanthan gum; dairy products such as milk; various kinds of fruit juices; proteins such as soybean protein; various kinds of phosphates; etc.

[0013] Yolk may be used in the acidic oil-in-water type emulsion composition according to the present invention for the purpose of emulsifying and improving flavor. Yolk may be used in any form such as raw, frozen, powdery, salted or sugared. Furthermore, it may also be used in the form of the whole egg containing egg white. The content of the yolk in the acidic oil-in-water type emulsion composition according to the present invention is preferably 5 to 20 %, more preferably 7 to 17 %, most preferably 8 to 15 % based on weight of raw yolk, from the viewpoint of improvement in flavor.

[0014] In the acidic oil-in-water type emulsion composition according to the present invention, a proportion (hereafter referred to as "lyso proportion") of lysophospholipid to total phospholipids contained is preferably at least 15 %, more preferably at least 25 %, most preferably 29 to 60 % from the viewpoints of shelf stability, appearance and flavor. A part or the whole of the lysophospholipid is preferably derived from yolk and/or soybean, most preferably from yolk. Furthermore, a part or the whole of the lysophospholipid is preferably enzyme-treated yolk. The enzyme used in the enzyme treatment of yolk is preferably esterase, lipase or phospholipase, more preferably lipase or phospholipase, most preferably phospholipase. Among various kinds of phospholipases, phospholipase A, i.e., phospholipase A1 or A2 is most preferred.

[0015] The acidic oil-in-water type emulsion composition according to the present invention preferably has a pH of 2.5 to 5.0, more preferably 3.0 to 4.5, most preferably 3.4 to 4.2.

[0016] The acidic oil-in-water type emulsion composition according to the present invention can be produced in accordance with, for example, the following process. Oil components such as DAG and the crystallization inhibitors are first mixed to prepare an oil phase. Yolk and other water-soluble raw materials are then mixed to prepare a water phase. The oil phase is added to the water phase, and the mixture is homogenized following preliminary emulsification if needed, whereby an acidic oil-in-water type emulsion composition can be obtained. Examples of homogenizers include high-pressure homogenizers such as Mountaingorin and Microfluidizer, ultrasonic emulsifiers, colloid mills, agitating homomixers, and Milder.

[0017] The average droplet diameter of oil droplets in the acidic oil-in-water type emulsion composition produced in such a manner is preferably 0.5 to 40.0 μm , more preferably 1.0 to 15.0 μm , most preferably 1.5 to 12.0 μm when measured by a laser diffraction method (using a laser diffraction particle size analyzer).

[0018] When the acidic oil-in-water type emulsion composition according to the present invention is mayonnaise or dressing, it is more preferable to have the following average droplet diameter. In the case of mayonnaise, the average droplet diameter is preferably 0.5 to 20.0 μm , more preferably 1.0 to 15.0 μm , most preferably 1.5 to 10.0 μm . In the case of dressing, the average droplet diameter is preferably 0.5 to 40.0 μm , more preferably 1.0 to 40.0 μm , most preferably 5.0 to 40.0 μm .

[0019] Examples of product form of the acidic oil-in-water type emulsion composition according to the present invention include those defined by JAS as dressing, semi-solid dressing, emulsified liquid dressing, mayonnaise, salad dressing and French dressing, but not limited to. All kinds of products, widely accepted as mayonnaise and dressing,

correspond thereto.

[0020] In the following examples, the HLB of the crystallization inhibitors was calculated out in accordance with the Griffin's empirical equation for polyglycerol fatty acid esters, and sorbitan fatty acid esters. For sucrose fatty acid esters, emulsification method was used to determine HLB.

Referential Example 1:

Preparation of Oil Composition 1

[0021] Lipozyme IM (product of Novo Nordisk Industry Co.) was added to a mixture of rapeseed oil fatty acid (650 parts by weight) and glycerol (107 parts by weight) and subjected to esterification reaction at 40°C for 5 hours under 7 hPa. The resultant reaction mixture was then subjected to molecular distillation (235°C, 0.07 hPa). The distillate thus obtained was then bleached, washed with water and deodorized at 235°C for 2 hours to obtain Oil Composition 1 having a composition shown in Table 1. Referential Example 2:

Preparation of Oil Composition 2

[0022] A mixture of soybean oil fatty acid (650 parts by weight), after the content of saturated fatty acids had been reduced by winterization, glycerol (107 parts by weight) and calcium hydroxide (2 parts by weight) was reacted at 230°C for 0.5 hours under nitrogen gas atmosphere, the reaction mixture was allowed to stand for 12 hours to separate the glycerol phase, and then the resultant oil phase (oil composition) was washed with a 50 % (by weight) aqueous solution of citric acid whose proportion was 2 parts by weight to 100 parts by weight of the oil phase followed by centrifugation to afford oil composition. The resultant oil composition was then subjected to molecular distillation (235°C, 0.07 hPa), and the distillate thus obtained was then bleached, washed with water and deodorized at 235°C for 2 hours to obtain Oil Composition 2 having a composition shown in Table 1.

Table 1

		Oil composition	
		1	2
Composition of *1oil	Triacylglycerol	13.8	13.5
	Diacylglycerol	84.7	85.2
	Monoacylglycerol	1.2	1.0
	Free fatty acid	0.3	0.3
Fatty acid *2composition	C16: 0	4.2	2.6
	C18: 0	1.9	0.7
	C18: 1	58.3	30.0
	C18: 2	21.3	57.5
	C18: 3	10.7	6.8
	C20: 0	0.7	1.5
	C20: 1	1.8	0.4
	C22: 0	0.2	0.1
	C22: 1	0.8	0.1

*1: Analyzed by gas chromatography after silylation.

*2: Analyzed by gas chromatography after methylation (Number of carbon atoms in fatty acid: number of carbon-carbon double bonds).

Referential Example 3:

Preparation 1 of enzyme-treated yolk

[0023] A yolk solution (750 g) containing common salt (10 % by weight) was mixed with water (250 g). After the resultant mixture was preheated at 50°C for 10 minutes, Phospholipase A2 (enzymatic activity: 10,000 IU/mL) was

added to the mixture. The proportion of phospholipase to yolk solution was 100 ppm. Reaction was carried out for 3 to 5 hours, thereby obtaining Enzyme-treated Yolk Solution 1.

[0024] After a yolk solution (850 g) containing common salt (10 % by weight) was mixed with water (150 g), and the resultant mixture was heated to 50°C, the above-described Phospholipase A2 was added to the mixture. The proportion of phospholipase to the yolk solution was 40 ppm. Reaction was carried out for 20 hours, thereby obtaining Enzyme-treated Yolk Solution 2.

[0025] After a yolk solution (850 g) containing common salt at a concentration of 10 % by weight was mixed with water (150 g), and the resultant mixture was preheated to 50°C, Phospholipase A1 (enzymatic activity: 4,000 IU/mL) was added to the mixture. The proportion of phospholipase to the yolk solution was 200 ppm. Reaction was carried out for 5 hours, thereby obtaining Enzyme-treated Yolk Solution 3.

[0026] The lyso proportion was calculated out in accordance with the following method.

[0027] The reaction product was extracted repeatedly with a mixed solvent (3:1) of chloroform:methanol to obtain a lipid mixture. The resultant lipid mixture was subjected to thin layer chromatography to separate various kinds of phospholipids by two-dimensional thin layer chromatography of the first dimension: chloroform-methanol-water (65:25:49) and the second dimension: butanol-acetic acid-water (60:20:20). The amounts of the separated phospholipids were measured by means of a commercially available measuring kit (permanganate ashing method, Phospholipid Test Wako, trade name; product of Wako Pure Chemical Industries, Ltd.) to calculate out contents of lysophospholipids and total phospholipids. The lyso proportion (%) was defined as "(total amount of phosphorus in lysophospholipid fraction/total amount of phosphorus in all phospholipid fractions) x 100".

Referential Example 4:

Preparation 2 of enzyme-treated yolk

[0028] Twice amount of water by volume was added to Enzyme-treated Yolk Solution 1 obtained in Referential Example 3 and stirred. The mixture was then spray-dried under air (inlet air temperature=170°C, outlet air temperature=70°C) to obtain enzyme-treated yolk powder. Examples 1 to 13 and Comparative Examples 1 to 6:

[0029] Oil phases and water phases of their corresponding compositions shown in Tables 2 and 3 were prepared in a conventional manner. After each of the oil phases was added to its corresponding water phase while stirring the water phase to preliminarily emulsify the oil phase, the emulsion was homogenized by a colloid mill (5,000 rpm, clearance: 0.35 mm) to prepare mayonnaise having an average emulsified droplet diameter of 2.5 to 3.5 µm. The mayonnaise was charged into a mayonnaise tube (100 g). In respect to Example 11 and Comparative Example 4, each mayonnaise was prepared having the average droplet diameter of 8.1 to 9.5 and then charged into a mayonnaise tube (100g)

[0030] Mayonnaise samples thus obtained were stored for 1 month to test under low temperature (-5, 0 and 5°C) and programmed temperature (max 25°C, min 0°C) and then left to stand at room temperature for 3 hours. Thereafter, the appearances and physical properties thereof were evaluated by 6 panels in accordance with the following evaluation standards. The average values of these evaluations are shown in Table 2.

Evaluation standards:

(1) Appearance:

[0031] The appearance of each mayonnaise sample charged into a tube was visually evaluated.

- 3: Good;
- 2: Signs of separation was observed;
- 1: Obvious separation has taken place.

(2) Physical property

[0032] The mayonnaise sample squeezed out of the tube was visually evaluated

- 3: Good;
- 2: Signs of roughening on the surface and/or water separation was observed;
- 1: Obvious roughening on surface and/or water separation has taken place.

Table 2

(% by weight)

		Example												
		1	2	3	4	5	6	7	8	9	10	11	12	13
Water phase	Common salt	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	3.0	1.2	1.8
	Sucrose	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	6.0	1.5
	Condiment	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	-	0.5
	Mustard powder	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.3
	Thickener ¹	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	-	0.2	0.6	-	0.6
	Thickener ²	-	-	-	-	-	-	-	-	0.2	-	-	-	-
	Thickener ³	-	-	-	-	-	-	-	-	-	-	-	1.5	2.0
	Soybean protein	-	-	-	-	-	-	-	-	-	-	1.5	-	-
	Enzyme-treated yolk 1	20.0	20.0	20.0	20.0	20.0	20.0	-	-	20.0	-	-	-	-
	Enzyme-treated yolk 2	-	-	-	-	-	-	-	-	-	-	-	7.0	-
	Enzyme-treated yolk 3	-	-	-	-	-	-	-	-	-	-	-	-	20.0
	Enzyme-treated yolk powder	-	-	-	-	-	-	-	-	-	-	-	-	-
	Yolk	-	-	-	-	-	-	7.4	3.7	-	15.0	-	-	-
Oil phase	10% Brewed vinegar	6.0	6.0	6.0	6.0	6.0	6.0	6.0	6.0	6.0	6.0	9.0	9.5	9.0
	Water	1.9	1.9	1.9	1.9	1.9	1.9	14.5	10.7	1.9	6.9	28.1	29.6	33.3
	Oil Composition 1	20.7	20.5	19.5	20.5	20.5	34.5	20.5	20.5	20.5	20.5	16.5	13.2	9.0
	Oil Composition 2	48.7	48.5	47.5	48.5	48.5	34.5	48.5	48.5	48.5	48.5	38.5	30.8	21.0
	Polyglycerol fatty acid ester ⁴	0.6	1.0	3.0	-	-	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	Sorbitan fatty acid ester ⁵	-	-	-	1.0	-	-	-	-	-	-	-	-	-
	Sucrose fatty acid ester ⁶	-	-	-	-	-	-	-	-	-	-	-	-	-
	Lysopropion (%)	45	45	45	45	45	50	55	55	50	-	-	55	50
	Appearance	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
	Physical property	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
	Appearance	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
	Physical property	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
	Appearance	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
Evaluation	Physical property	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
	Appearance	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
	Physical property	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
	Appearance	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0
	Physical property	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0

Table 3

		Comparative Example						(% by weight)
		1	2	3	4	5	6	
Water phase	Common salt	0.3	0.3	0.3	3.0	1.2	1.8	
	Sucrose	1.0	1.0	1.0	1.0	6.0	1.5	
	Condiment	0.4	0.4	0.4	0.5	-	0.5	
	Mustard powder	0.2	0.2	0.2	0.3	0.2	0.3	
	Thickener *1	0.2	0.2	0.2	0.6	-	0.6	
	Thickener *2	-	-	-	-	-	-	
	Thickener *3	-	-	-	1.5	1.5	2.0	
	Soybean protein	-	-	-	-	-	-	
	Enzyme-treated yolk 1	20.0	20.0	20.0	-	-	-	
	Enzyme-treated yolk 2	-	-	-	-	7.0	-	
Oil phase	Enzyme-treated yolk 3	-	-	-	-	-	20.0	
	Enzyme-treated yolk powder	-	-	-	-	-	-	
	Yolk	-	-	-	-	-	-	
	10% Brewed vinegar	6.0	6.0	6.0	9.0	9.5	9.0	
	Water	1.9	1.9	1.9	29.1	30.6	34.3	
	Oil Composition 1	21.0	35.0	20.8	16.5	13.2	9.0	
	Oil Composition 2	49.0	35.0	48.8	38.5	30.8	21.0	
	Polyglycerol fatty acid ester *4	-	-	0.4	-	-	-	
	Sorbitan fatty acid ester *5	-	-	-	-	-	-	
	Sucrose fatty acid ester *6	-	-	-	-	-	-	
Lyso proportion (%)		45	50	50	-	55	50	
Evaluation	Appearance	3.0	3.0	3.0	3.0	3.0	3.0	
	Physical property	3.0	3.0	3.0	3.0	3.0	3.0	
	Appearance	3.0	2.0	3.0	3.0	3.0	3.0	
	Physical property	2.5	1.5	3.0	3.0	3.0	3.0	
	Appearance	1.0	1.0	1.0	2.0	2.5	2.7	
	Physical property	1.0	1.0	1.0	2.0	2.5	2.7	
	Appearance	3.0	3.0	3.0	3.0	3.0	3.0	
	Physical property	3.0	3.0	3.0	3.0	3.0	3.0	
	Cycle test	3.0	3.0	3.0	3.0	3.0	3.0	
	Physical property	3.0	3.0	3.0	3.0	3.0	3.0	

*1: Xanthan gum (product of DAINIPPON PHARMACEUTICAL CO., LTD).

*2: Tamarind gum (product of DAINIPPON PHARMACEUTICAL CO., LTD).

*3: INSTANT PURE-FLO (product of National Starch and Chemical Company).

*4: Average polymerization degree: 10; degree of esterification: at least 80%; fatty acid composition: C18 and C22:1; HLB: 2.0.

*5: Fatty acid composition: C18; HLB: 2.1.

*6: Fatty acid composition: C18 and C2; degree of esterification of C18: at least 60%; acetylation rate: at least 35%; HLB: 0.

[0033] Comparative Examples 1 and 2 relate to mayonnaise devoid of crystallization inhibitors, and both mayonnaise samples underwent demulsification (oil-off) after storage for 1 month at -5°C and 0°C and caused separation, surface roughness and water separation. Comparative Example 3 relates to mayonnaise in which excessive amount of polyglycerol fatty acid ester (0.4 % by weight) was used in the oil phase. Sufficient emulsion stability was unable to be achieved in this mayonnaise

[0034] On the other hand, when a proper amount of polyglycerol fatty acid ester (Examples 1 to 3 and 6 to 8), sorbitan fatty acid ester (Example 4) or sucrose fatty acid ester (Example 5) was used as crystallization inhibitor, no demulsification was observed even after storage for 1 month at -5°C to 5°C, and both appearance and physical property were good.

[0035] Emulsion stability tends to be improved when the ratio of oil phase is lowered, but sufficient improvement cannot be obtained in the absence of crystallization inhibitor. When used in a proper amount, the crystallization inhibitor was able to bring sufficient improvement in the emulsion stability (Examples 11 to 13 and Comparative Examples 4 to 6).

[0036] Furthermore, the polyglycerol fatty acid ester (Examples 1 to 3 and 6 to 8) was confirmed to be the best use as a crystallization inhibitor agent both in appearance and physical property, from results of the tests under programmed temperature.

Examples 14 to 16:

[0037] French dressing (Example 14), Thousand Island dressing (Example 15) and sesame dressing (Example 16) were produced with their corresponding compositions shown in Table 4. More specifically, raw materials for an oil phase were added dropwise to a water phase under stirring to conduct preliminary emulsification. This emulsion was homogenized by a homomixer to obtain the respective dressing samples having an average emulsified droplet diameter of 4 to 12 µm. With respect to the respective dressing samples, the same evaluation as in Example 1 was conducted by 6 panelists. As a result, all the dressing samples were good in both appearance and physical property even after storage for 1 month at -5°C to 5°C as shown in Table 4.

Table 4

(% by weight)

		Example			
		14	15	16	
Water phase	Common salt	3.0	2.0	2.5	
	Sucrose	5.0	5.0	11.0	
	Condiment	0.5	0.5	1.0	
	Lemmon juice	2.0	2.0	-	
	Thickener ^{*1}	0.6	0.4	0.01	
	Enzyme-treated yolk 1	2.0	4.0	-	
	Yolk	-	-	1.5	
	Tomato ketchup	-	5.0	-	
	Pickles	-	4.0	-	
	Tomato paste	-	1.0	-	
	Miso	-	-	4.0	
	Pounded sesame	-	-	7.0	
	Soy	-	-	3.0	
	5% Brewed vinegar	14.0	14.0	14.0	
Water	32.9	27.1	25.99		
Oil phase	Oil Composition 1	19.5	17.0	14.5	
	Oil Composition 2	19.5	17.0	14.5	
	Polyglycerol fatty acid ester ^{*2}	1.0	1.00	1.0	
Lyso proportion (%)		40	40	-	
Evaluation	5°C	Appearance	3.0	3.0	3.0
		Physical property	3.0	3.0	3.0
	0°C	Appearance	3.0	3.0	3.0
		Physical property	3.0	3.0	3.0
	-5°C	Appearance	3.0	3.0	3.0
		Physical property	3.0	3.0	3.0

*1: Xanthan gum (product of DAINIPPON PHARMACEUTICAL CO., LTD).

*2: Average polymerization degree: 10; degree of esterification: at least 80%; fatty acid composition: C18: 1 and C16.

Industrial Applicability

[0038] As described above, the acidic oil-in-water type emulsion compositions according to the present invention are so excellent in shelf stability at low temperatures that they do not crack during storage at a low temperature in a refrigerator though they contain DAG at a high concentration, also good in appearance and flavor and useful as diets or lipid metabolism-improving foods.

Claims

1. An acidic oil-in-water type emulsion composition comprising an oil phase containing at least 20 % by weight of diacylglycerol and 0.5 to 5.0 % by weight of crystallization inhibitor.

EP 1 214 886 A1

2. The acidic oil-in-water type emulsion composition according to Claim 1, wherein the crystallization inhibitor is selected from polyglycerol fatty acid esters, sucrose fatty acid esters and sorbitan fatty acid esters.
- 5 3. The acidic oil-in-water type emulsion composition according to Claim 2, wherein the polyglycerol fatty acid esters are such that the average polymerization degree of glycerol is 2 to 12, the number of carbon atoms in the fatty acid moiety is 12 to 22, and the degree of esterification is at least 70 %.
- 10 4. The acidic oil-in-water type emulsion composition according to Claim 2, wherein the sucrose fatty acid esters are such that the degree of esterification with a fatty acid having 12 to 22 carbon atoms is at least 50 %, and the remaining hydroxyl groups (not esterified with above-mentioned fatty acids) are acetylated.
- 15 5. The acidic oil-in-water type emulsion composition according to Claim 2, wherein the sorbitan fatty acid esters are those in which the number of carbon atoms in the fatty acid composition is 12 to 22, and the HLB is lower than 3.
6. The acidic oil-in-water type emulsion composition according to any one of Claims 1 to 5, which further comprises egg yolk.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 01 12 9299

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
P,Y	WO 00 78162 A (KONISHI YOSHIHIRO ;KAWAI SHIGERU (JP); KAO CORP (JP)) 28 December 2000 (2000-12-28) * claims 1,7 *	1,2,5,6	A23D7/00 A61K7/48 A23L1/24 A61K47/14
P,Y	EP 1 135 991 A (KAO CORP) 26 September 2001 (2001-09-26) * page 4, line 32-45; claims 1,4,6 *	1,2,5,6	
Y	EP 0 378 893 A (KAO CORP) 25 July 1990 (1990-07-25) * page 6, line 8-12; claim 1 *	1-3,5,6	
X	EP 0 836 805 A (KAO CORP) 22 April 1998 (1998-04-22) * claims 1,7,10 *	1,2 3-6	
Y	EP 0 402 090 A (KAO CORP) 12 December 1990 (1990-12-12) * page 6, line 37-42; claim 1 *	1-3,5,6	
Y	EP 0 750 848 A (KAO CORP) 2 January 1997 (1997-01-02) * claims 32,34 *	1,2	TECHNICAL FIELDS SEARCHED (Int.Cl.7) A23D A61K A23L
Y	EP 0 990 391 A (KAO CORP) 5 April 2000 (2000-04-05) * claims 1,6,9,10 *	1,2,6	
Y	US 5 340 600 A (RECKWEG FREEK ET AL) 23 August 1994 (1994-08-23) * column 2, line 13-22 *	1-3,5,6	
-/--			
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 16 April 2002	Examiner Groh, B
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPFO FORM 1503.03.02 (P01001)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 01 12 9299

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y	<p>DATABASE WPI Section Ch, Week 198742 Derwent Publications Ltd., London, GB; Class D13, AN 1987-294789 XP002196090 & JP 62 205738 A (MITSUBISHI CHEM IND LTD) , 10 September 1987 (1987-09-10) 'Title: "Agent for suppressing crystallization of oil or fat - comprises sucrose palmitate of HLB below 5"! * abstract *</p>	1-6	
Y	<p>US 5 660 865 A (PEDERSEN ARNE ET AL) 26 August 1997 (1997-08-26) * column 3, line 29-31 *</p>	1-6	
Y	<p>DATABASE WPI Section Ch, Week 199807 Derwent Publications Ltd., London, GB; Class D13, AN 1998-072028 XP002196091 & JP 09 310088 A (RIKEN VITAMIN CO), 2 December 1997 (1997-12-02) 'Title: Crystallisation inhibitor for liquid edible fat and oil - comprises mixture of poly-glycerine fatty acid ester and sorbitan fatty acid ester! * abstract *</p>	1-6	
A	<p>DATABASE WPI Section Ch, Week 199218 Derwent Publications Ltd., London, GB; Class D13, AN 1992-142505 XP002196208 "Nutritional oil-in-water emulsion" & JP 04 264033 A, 18 September 1992 (1992-09-18) * abstract *</p>	1-6	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 16 April 2002	Examiner Groh, B
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons 5 : member of the same patent family, corresponding document</p>			

EPO FORM 503 03.82 (PO/C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 12 9299

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

16-04-2002

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0078162	A	28-12-2000	JP	2001000138 A	09-01-2001
			BR	0012257 A	12-03-2002
			EP	1185179 A2	13-03-2002
			WO	0078162 A2	28-12-2000
EP 1135991	A	26-09-2001	JP	2001262180 A	26-09-2001
			EP	1135991 A2	26-09-2001
			US	2001036502 A1	01-11-2001
EP 0378893	A	25-07-1990	JP	2010558 C	02-02-1996
			JP	2190146 A	26-07-1990
			JP	7038771 B	01-05-1995
			DE	68907477 D1	12-08-1993
			DE	68907477 T2	21-10-1993
			DE	68923977 D1	28-09-1995
			DE	68923977 T2	01-02-1996
			EP	0378893 A2	25-07-1990
			EP	0525915 A1	03-02-1993
			ES	2041989 T3	01-12-1993
			ES	2076667 T3	01-11-1995
			HK	49894 A	27-05-1994
			PH	26001 A	29-01-1992
			SG	59594 G	28-10-1994
			US	4976984 A	11-12-1990
EP 0836805	A	22-04-1998	CN	1181194 A , B	13-05-1998
			EP	0836805 A1	22-04-1998
			JP	10176181 A	30-06-1998
			US	6004611 A	21-12-1999
EP 0402090	A	12-12-1990	JP	2848849 B2	20-01-1999
			JP	3008431 A	16-01-1991
			JP	2777217 B2	16-07-1998
			JP	3089936 A	15-04-1991
			JP	2704004 B2	26-01-1998
			JP	3091451 A	17-04-1991
			DE	69003014 D1	07-10-1993
			DE	69003014 T2	16-12-1993
			EP	0402090 A2	12-12-1990
			ES	2043279 T3	16-12-1993
			HK	56494 A	03-06-1994
			US	5160759 A	03-11-1992
EP 0750848	A	02-01-1997	JP	3017035 B2	06-03-2000
			JP	7228550 A	29-08-1995
			BR	9506809 A	14-10-1997

EPO FORM PUA/93

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 12 9299

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

16-04-2002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0750848	A		EP 0750848 A1	02-01-1997
			US 5789011 A	04-08-1998
			WO 9619118 A1	27-06-1996
EP 0990391	A	05-04-2000	BR 9904796 A	30-05-2000
			EP 0990391 A1	05-04-2000
			CN 1258199 T	28-06-2000
			WO 9948378 A1	30-09-1999
			US 6139897 A	31-10-2000
			US 6326050 B1	04-12-2001
US 5340600	A	23-08-1994	AT 151951 T	15-05-1997
			AU 684490 B2	18-12-1997
			AU 6000694 A	14-09-1994
			CA 2156265 A1	01-09-1994
			CZ 9502099 A3	13-03-1996
			DE 69402831 D1	28-05-1997
			DK 684769 T3	27-10-1997
			WO 9418846 A1	01-09-1994
			EP 0684769 A1	06-12-1995
			PL 310170 A1	27-11-1995
			SK 100495 A3	06-12-1995
			ZA 9400993 A	14-08-1995
JP 62205738	A	10-09-1987	JP 1993740 C	22-11-1995
			JP 7024547 B	22-03-1995
US 5660865	A	26-08-1997	AT 153832 T	15-06-1997
			AU 674349 B2	19-12-1996
			AU 5107793 A	26-04-1994
			BR 9307109 A	30-03-1999
			CA 2145205 A1	14-04-1994
			DE 69311352 D1	10-07-1997
			DE 69311352 T2	02-01-1998
			WO 9407373 A1	14-04-1994
			DK 661924 T3	14-07-1997
			EP 0661924 A1	12-07-1995
			ES 2102680 T3	01-08-1997
			FI 951413 A	24-03-1995
			JP 8501454 T	20-02-1996
			NO 951120 A	18-05-1995
			NZ 256387 A	25-06-1996
JP 9310088	A	02-12-1997	NONE	
JP 4264033	A	18-09-1992	CH 681343 A5	15-03-1993

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 12 9299

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

16-04-2002

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 4264033 A	AT	114933 T	15-12-1994
	AU	635447 B2	18-03-1993
	AU	8484891 A	30-04-1992
	CA	2052988 A1	27-04-1992
	CN	1060948 A , B	13-05-1992
	DE	69105697 D1	19-01-1995
	DE	69105697 T2	13-04-1995
	DK	482345 T3	01-05-1995
	EP	0482345 A1	29-04-1992
	ES	2066305 T3	01-03-1995
	FI	914988 A	27-04-1992
	IE	913326 A1	22-05-1992
	JP	2023443 C	26-02-1996
	JP	7047544 B	24-05-1995
	NO	914164 A	27-04-1992
	NZ	240011 A	26-08-1993
	PT	99329 A	30-09-1992
	US	5192577 A	09-03-1993
	ZA	9107767 A	24-06-1992
<hr/>			

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82